

THE HEIGHT OF q -BINARY SEARCH TREES

MICHAEL DRMOTA AND HELMUT PRODINGER

ABSTRACT. q -binary search trees are obtained from words, equipped with the geometric distribution instead of permutations. The average and variance of the height are computed, based on random words of length n , as well as a Gaussian limit law.

1. INTRODUCTION

The paper [8] introduces for the first time a meaningful q -model of binary search trees: instead of binary search trees, one considers tournament trees, which differ only marginally from binary search trees; if one starts from a permutation $(\begin{smallmatrix} 1 & 2 & \dots & n \\ \pi_1 & \pi_2 & \dots & \pi_n \end{smallmatrix})$, then one inserts the number i instead of the number π_i . Thus, traversing the tree in inorder, we might think of the associated permutation as $\rho 1 \sigma$, where 1 goes into the root, and ρ resp. σ form (recursively) the left resp. right subtree. We could have called this paper “The Height of q -Tournament Trees;” however we decided not to do so since binary search trees are by far better known, both, in the community of theoretical computer scientists, and combinatorialists. A nice reference for tournament trees and increasing trees in general is [2].

Now instead of considering permutations $\pi_1 \pi_2 \dots \pi_n$, we consider words over the alphabet $\{1, 2, 3, \dots\}$, and (geometric) probabilities attached to the letters, i. e., the probability of letter i is pq^{i-1} , with $p+q = 1$. The binary search tree is then constructed by writing a nonempty word w as $w = xay$, where a is the smallest letter occurring, and $x \in \{a+1, a+2, \dots\}^*$ and $y \in \{a, a+1, \dots\}^*$. The letter a goes into the root and x resp. y form the left resp. right subtree.

The paper [8] dealt with the path length; here we consider the height. The height of a binary search tree (and thus of the trees in our model) is defined to be the largest number of nodes in a path from the root to a leaf; the empty tree (related to the empty word) has height 0.

We will prove that the expected height, when considering random words of length n , is asymptotic to pn ; the variance will also be computed as well as a Gaussian limit law. (The letter p will always denote $1 - q$ in this paper.) Recall that the result for traditional binary search trees is $\sim c \log n$ with $c = 4.31107$, see [3]; unfortunately we do not get that result as the limit $q \rightarrow 1$ as it so happened for the path length. However, nothing is wrong here, since the limit $q \rightarrow 1$ in pn simply tells us that the average height should be less than linear. Although this original hope of getting the

Key words and phrases. Binary search tree, q -analog, height.

classical result as a corollary did not work out, we nevertheless think that the results presented here are of independent interest. Note also that pn is the expected number of letters 1 in a (random) string of length n . In our asymmetric model, they must all lie on one path.

In more detail, we will obtain the following results:

Let H_n denote the height of q -binary search trees with n (internal) nodes (this is a random variable, defined on words of length n).

Theorem 1. *For every positive $q < 1$ the height H_n of q -binary search trees with n (internal) nodes satisfies a central limit theorem of the form*

$$\sup_{x \in \mathbb{R}} \left| \mathbf{Pr}\{H_n \leq x\} - \Phi\left(\frac{x - pn}{\sqrt{npq}}\right) \right| = \mathcal{O}(n^{-1/2}), \quad (1)$$

where $\Phi(x)$ denotes the normal distribution function

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt.$$

The expected value is given by

$$\mathbb{E} H_n = pn + \mathcal{O}\left(\frac{1}{p}\right), \quad (2)$$

in which the \mathcal{O} -constant is uniform for $0 < q < 1$. The variance can be estimated by

$$\mathbb{V} H_n = pqn + \mathcal{O}_q\left(n^{1/2} \log^2 n\right). \quad (3)$$

The \mathcal{O} -constant in the error term of the variance is not uniform for $0 < q < 1$. The dependency on q has not been worked out since the order of magnitude of the error term is surely not optimal. However, although the error term for the expected value is uniform, this theorem cannot be used to cover the case $q \rightarrow 1$ where it is known that the expected value is given by

$$\mathbb{E} H_n = c \log n + \mathcal{O}(\log \log n)$$

(with $c = 4.31107\dots$, see [3]) and the variance is bounded:

$$\mathbb{V} H_n = \mathcal{O}(1)$$

(see [5, 6]).

Intuitively, this result (and its proof) says that the height is dominated by the number of 1's in the sequence. This seems to be due to the fact that most 1's appear consecutively in the sequence, producing rather skew subtrees. The contribution from 2's is asymptotically negligible since there are much fewer of them, etc.

Studying the combinatorics (of words) of geometrically distributed random variables has been a long term project of one of us (H. P.), and further papers can be found on this author's webpage. We do not want to give more details here, but in all previous

cases the correspondence between the model of words and its limit (permutations) led to very satisfactory results.

We would like to cite the paper [1] which is of general interest in this context.

2. WEAK CONVERGENCE

Lemma 1. *Let $y_k(x)$ denote the generating function*

$$y_k(x) = \sum_{n \geq 0} \Pr\{H_n \leq k\} x^n.$$

Then we have $y_0(x) \equiv 0$, $y_k(0) = 1$ and

$$y_{k+1}(x) = px y_k(x) y_k(qx) + y_{k+1}(qx) \quad (4)$$

for $k \geq 0$.

Proof. The proof is a straightforward translation of the basic decomposition $w = xay$: If $a = 1$ (described by px), then the left subtree consists only of letters in $\{2, 3, \dots\}$, described by $y_k(qx)$, but the right subtree can have any letters, which is described by $y_k(x)$. However, if $a > 1$, we get the term $y_{k+1}(qx)$ (we might then think of all letters being reduced by 1). \square

If we write this in the form

$$\frac{y_{k+1}(x) - y_{k+1}(qx)}{(1-q)x} = y_k(qx) y_k(x),$$

then the limit $q \rightarrow 1$ gives

$$y'_{k+1}(x) = y_k^2(x),$$

the usual recursion in the instance of binary search trees. (Recall that $\frac{f(x)-f(qx)}{(1-q)x} =: (D_q f)(x)$, and D_q is called the q -difference operator.) Also, if we write

$$y_k(x) = \sum_{0 \leq j < 2^k} a_{k,j} x^j,$$

then we get, by comparing coefficients,

$$a_{k+1,j} = \frac{p}{1-q^j} \sum_{0 \leq i < j} a_{k,i} a_{k,j-1-i} q^i \quad \text{for } j \geq 1,$$

and $a_{k,0} = 1$ for all $k \geq 0$. Thus, Theorem 1 may be reformulated in terms of $a_{k,j}$. However, in this paper we will not make use of this notion.

Lemma 2. *The generating functions $y_k(x)$ ($k \geq 0$, $0 \leq x < 1$) are bounded above by*

$$y_k(x) \leq \frac{1}{1-x} - \frac{x}{1-x} \left(\frac{px}{1-qx} \right)^k. \quad (5)$$

Furthermore, this inequality is also true on the level of coefficients, i. e. for $n \geq 1$ and $k \geq 1$

$$\begin{aligned} \Pr\{H_n \leq k\} &= [x^n]y_k(x) \\ &\leq 1 - [x^n]\frac{x}{1-x} \left(\frac{px}{1-qx}\right)^k \\ &= \sum_{l=0}^{k-1} \binom{n-1}{l} p^l q^{n-1-l}. \end{aligned} \tag{6}$$

Note that this upper bound is an exact binomial distribution function which is asymptotically normal with mean $p(n-1) + 1 = pn + q$ and variance $pq(n-1)$.

Proof. Since $y_k(x) \leq 1/(1-x)$ we get from (4) that

$$y_{k+1}(x) \leq \frac{px}{1-qx}y_k(x) + \frac{1}{1-qx}.$$

Thus, (5) follows by induction. Note that any step in these calculations is also true on the level of coefficients. Thus,

$$\Pr\{H_n \leq k\} \leq 1 - [x^n]\frac{x}{1-x} \left(\frac{px}{1-qx}\right)^k.$$

Set

$$Y_k(x) = \frac{1}{1-x} - \frac{x}{1-x} \left(\frac{px}{1-qx}\right)^k$$

and

$$\Delta_k(x) = Y_k(x) - Y_{k-1}(x) = \frac{1}{p} \left(\frac{px}{1-qx}\right)^k.$$

Since

$$[x^n]\Delta_k(x) = \binom{n-1}{k-1} p^{k-1} q^{n-k},$$

(6) follows immediately. \square

Lemma 3. *The generating functions $y_k(x)$ ($k \geq 0$, $0 \leq x < 1$) are bounded below by*

$$\begin{aligned}
y_k(x) &\geq \frac{1}{1-x} - \frac{x}{1-x} \left(\frac{px}{1-qx} \right)^k \\
&\quad - \frac{(qx)^2}{1-qx} \frac{\left(\frac{px}{1-qx} \right)^k - (qx)^k}{\frac{px}{1-qx} - qx} \\
&\quad - \frac{qx}{1-x} \left(\frac{px}{1-qx} \right) \frac{\left(\frac{px}{1-qx} \right)^k - (qx)^k}{\frac{px}{1-qx} - qx} \\
&\quad + \frac{qx^2}{1-x} \left(\frac{px}{1-qx} \right)^k \frac{1 - (qx)^k}{1 - qx}.
\end{aligned} \tag{7}$$

Furthermore, this inequality is also true on the level of coefficients.

Proof. First we use the trivial lower bound

$$y_k(x) \geq \frac{1 - x^{k+1}}{1 - x}$$

and the upper bound (5) to obtain the inequality

$$\begin{aligned}
y_{k+1}(x) &= px y_k(x) y_k(qx) + y_{k+1}(qx) \\
&\geq x(1-q) y_k(x) \left(\frac{1}{1-qx} - \frac{(qx)^{k+1}}{1-qx} \right) + \frac{1}{1-qx} - \frac{(qx)^{k+2}}{1-qx} \\
&\geq \frac{px}{1-qx} y_k(x) + \frac{1}{1-qx} \\
&\quad - px \frac{(qx)^{k+1}}{1-qx} \left(\frac{1}{1-x} - \frac{x}{1-x} \left(\frac{px}{1-qx} \right)^k \right) - \frac{(qx)^{k+2}}{1-qx}.
\end{aligned}$$

Now (7) follows by induction. As in the proof of Lemma 2 it can be observed that (7) is also true on the level of coefficients. \square

In order to obtain proper error terms for the lower bound for $\Pr\{H_n \leq k\}$ which may derived from Lemma 3 we make use of the following lemma.

Lemma 4. *Let $0 < q < 1$ be given and let $F(x)$ be a function which is analytic for $|x| < 1 + \varepsilon$ for some $\varepsilon > 0$. Then there exist $c > 0$ and $\eta > 0$ (depending on q) such that*

$$[x^n]F(x) \left(\frac{px}{1-qx} \right)^k = \mathcal{O} \left(n^{-1/2} \exp \left(-\frac{c}{n} (k - pn)^2 \right) \right) \tag{8}$$

and

$$[x^n]F(x) \frac{\left(\frac{px}{1-qx} \right)^k - (qx)^k}{\frac{px}{1-qx} - qx} = \mathcal{O} \left(n^{-1/2} \exp \left(-\frac{c}{n} (k - pn)^2 \right) \right) + \mathcal{O}(q^k) \tag{9}$$

uniformly for $|k - pn| \leq \eta n$ as $n \rightarrow \infty$.

Proof. By using standard saddle point asymptotics (compare with [4]) it follows that

$$\begin{aligned} [x^n]F(x) \left(\frac{px}{1-qx} \right)^k &= \frac{1}{2\pi i} \int_{|z|=x_0} F(z) \left(\frac{pz}{1-qz} \right)^k \frac{dz}{z^{n+1}} \\ &= \mathcal{O} \left(n^{-1/2} x_0^{-n} \left(\frac{px_0}{1-qx_0} \right)^k \right), \end{aligned}$$

where the saddle point $x = x_0 = \left(1 - \frac{k}{n}\right)/q$ is determined by the equation

$$\frac{x \frac{d}{dx} \left(\frac{px}{1-qx} \right)}{\frac{px}{1-qx}} = \frac{n}{k}.$$

In particular we have $x_0 = 1$ if $k/n = p$. Finally a local expansion of

$$x_0^{-n} \left(\frac{px_0}{1-qx_0} \right)^k = \frac{q^{n-k} p^k}{\left(1 - \frac{k}{n}\right)^{n-k} \left(\frac{k}{n}\right)^k}$$

completes the proof of (8).

The proof of (9) runs along similar lines. However, we have to be a little bit more careful. The reason is that the denominator

$$\frac{pz}{1-qz} - qz$$

is singular for $z_1 = 0$ and for $z_2 = (-1 + 2q)/q^2$. Nevertheless the whole part

$$\frac{\left(\frac{pz}{1-qz} \right)^k - (qz)^k}{\frac{pz}{1-qz} - qz}$$

is regular for $|z| < 1/q$. Note that $|z_2| > 1$ for $0 < q < \sqrt{2} - 1$ and that $|z_2| < 1$ for $\sqrt{2} - 1 < q < 1$. Thus, if $q \neq \sqrt{2} - 1$ we get similarly to the above

$$\begin{aligned}
[x^n]F(x) \frac{\left(\frac{px}{1-qx}\right)^k - (qx)^k}{\frac{px}{1-qx} - qx} &= \frac{1}{2\pi i} \int_{|z|=x_0} F(z) \frac{\left(\frac{pz}{1-qz}\right)^k - (qz)^k}{\frac{pz}{1-qz} - qz} \frac{dz}{z^{n+1}} \\
&= \frac{1}{2\pi i} \int_{|z|=x_0} F(z) \frac{\left(\frac{pz}{1-qz}\right)^k}{\frac{pz}{1-qz} - qz} \frac{dz}{z^{n+1}} - \frac{1}{2\pi i} \int_{|z|=x_0} F(z) \frac{(qz)^k}{\frac{pz}{1-qz} - qz} \frac{dz}{z^{n+1}} \\
&= \frac{1}{2\pi i} \int_{|z|=x_0} F(z) \frac{\left(\frac{pz}{1-qz}\right)^k}{\frac{pz}{1-qz} - qz} \frac{dz}{z^{n+1}} - \frac{1}{2\pi i} \int_{|z|=1} F(z) \frac{(qz)^k}{\frac{pz}{1-qz} - qz} \frac{dz}{z^{n+1}} \\
&= \mathcal{O}\left(n^{-1/2} x_0^{-n} \left(\frac{px_0}{1-qx_0}\right)^k\right) + \mathcal{O}(q^k).
\end{aligned}$$

Note that we just shifted the paths of integration in regions of analyticity if η is chosen sufficiently small such that $|x_0 - 1| < |1 - z_2|$.

If $q = \sqrt{2} - 1$ then there are polar singularities at $z_2 = -1$ after splitting the integral. However, the residues of the functions involved (at $z = -1$) are both of order

$$\mathcal{O}\left((\sqrt{2} - 1)^k\right) = \mathcal{O}(q^k),$$

which implies that we get the same error term as in the case $q \neq \sqrt{2} - 1$. \square

Note further that (with a little bit more effort) we could have been much more precise. However, the bound given by Lemma 4 is sufficient for our purposes.

Lemma 5. *For every $0 < q < 1$ there exist $c > 0$ and $\eta > 0$ (depending on q) such that*

$$\left| \Pr\{H_n \leq k\} - \sum_{l=0}^k \binom{n-1}{l} p^l q^{n-1-l} \right| = \mathcal{O}\left(n^{-1/2} \exp\left(-\frac{c}{n}(k-pn)^2\right)\right) \quad (10)$$

uniformly for $|k - pn| \leq \eta n$ and

$$\left| \Pr\{|H_n - pn| \geq m\} \right| = \mathcal{O}\left(\exp\left(-c\frac{m^2}{n}\right)\right) \quad (11)$$

uniformly for $|m| \leq \eta n$ as $n \rightarrow \infty$.

Proof. First we note that (9) of Lemma 4 applies to the function

$$\frac{(qx)^2}{1-qx} \frac{\left(\frac{px}{1-qx}\right)^k - (qx)^k}{\frac{px}{1-qx} - qx}.$$

Thus we obtain

$$[x^n] \frac{(qx)^2}{1-qx} \frac{\left(\frac{px}{1-qx}\right)^k - (qx)^k}{\frac{px}{1-qx} - qx} = \mathcal{O}\left(n^{-1/2} \exp\left(-\frac{c}{n}(k-pn)^2\right)\right) + \mathcal{O}(q^k)$$

uniformly for $|k-pn| \leq \eta n$ as $n \rightarrow \infty$.

The remaining terms will now be treated in the same way as in the proof of Lemma 4:

$$\begin{aligned} & [x^n] \left(\frac{qx}{1-x} \left(\frac{px}{1-qx} \right) \frac{\left(\frac{px}{1-qx}\right)^k - (qx)^k}{\frac{px}{1-qx} - qx} + \frac{qx^2}{1-x} \left(\frac{px}{1-qx} \right)^k \frac{1 - (qx)^k}{1-qx} \right) \\ &= \frac{1}{2\pi i} \int_{|z|=1-\varepsilon} \left(\frac{qz}{1-z} \left(\frac{pz}{1-qz} \right) \frac{\left(\frac{pz}{1-qz}\right)^k - (qz)^k}{\frac{pz}{1-qz} - qz} - \frac{qz^2}{1-z} \left(\frac{pz}{1-qz} \right)^k \frac{1 - (qz)^k}{1-qz} \right) \frac{dz}{z^{n+1}} \\ &= \frac{1}{2\pi i} \int_{|z|=x_0} \left(\frac{qz}{1-z} \left(\frac{pz}{1-qz} \right) \frac{1}{\frac{pz}{1-qz} - qz} - \frac{qz^2}{1-z} \frac{1}{1-qz} \right) \left(\frac{pz}{1-qz} \right)^k \frac{dz}{z^{n+1}} \\ &\quad - \frac{1}{2\pi i} \int_{|z|=1-k^{-1}} \frac{qz}{1-z} \left(\frac{pz}{1-qz} \right) \frac{(qz)^k}{\frac{pz}{1-qz} - qz} \frac{dz}{z^{n+1}} \\ &\quad + \frac{1}{2\pi i} \int_{|z|=1-k^{-1}} \frac{qz^2}{1-z} \left(\frac{pz}{1-qz} \right)^k \frac{(qz)^k}{1-qz} \frac{dz}{z^{n+1}} \\ &= \mathcal{O}\left(n^{-1/2} x_0^{-n} \left(\frac{px_0}{1-qx_0} \right)^k\right) + \mathcal{O}(kq^k). \end{aligned}$$

For the first integral we use the fact that the function

$$F(z) = \frac{qz}{1-z} \left(\frac{pz}{1-qz} \right) \frac{1}{\frac{pz}{1-qz} - qz} - \frac{qz^2}{1-z} \frac{1}{1-qz}$$

is regular at $z = 1$ und thus bounded in a vicinity of $z = 1$ (compare with the proof of Lemma 4). The second and third integrals are easy to estimate. We only have to apply the trivial bounds

$$\max_{|z|=1-k^{-1}} \left| \frac{1}{1-z} \right| = k,$$

$$\max_{|z|=1-k^{-1}} |z^k| = \mathcal{O}(1)$$

and

$$\max_{|z|=1-k^{-1}} \left| \frac{pz}{1-qz} \right|^k = \left(1 + \mathcal{O}(k^{-1})\right)^k = \mathcal{O}(1).$$

Finally observe that in the range $|k - pn| \leq \eta n$ we surely have

$$kq^k = \mathcal{O}\left(n^{-1/2} \exp\left(-\frac{c}{n}(k - pn)^2\right)\right)$$

which completes the proof of (10).

The proof of (11) is now easy. We just have to combine a proper tail estimate for the binomial distribution, i. e.

$$\sum_{|l-pn| \geq m} \binom{n-1}{l} p^l q^{n-1-l} = \mathcal{O}\left(\exp\left(-c\frac{m^2}{n}\right)\right),$$

with (10). \square

We are now able to complete the first part of Theorem 1. It is well known that the distribution function of the binomial distribution can be estimated by the distribution function of the normal distribution up to a uniform error of order $\mathcal{O}(n^{-1/2})$, i. e., as $n \rightarrow \infty$,

$$\sup_{x \in \mathbb{R}} \left| \sum_{l \leq x} \binom{n-1}{l} p^l q^{n-1-l} - \Phi\left(\frac{x - pn}{\sqrt{pqn}}\right) \right| = \mathcal{O}(n^{-1/2}), \quad (12)$$

compare with [7, p. 542].

Furthermore, by (10) we get a similar result for the distribution of H_n :

$$\sup_{x \in \mathbb{R}} \left| \Pr\{H_n \leq x\} - \sum_{l \leq x} \binom{n-1}{l} p^l q^{n-1-l} \right| = \mathcal{O}(n^{-1/2}). \quad (13)$$

Note that (10) provides (13) just for x with $|x - pn| \leq \eta n$. However, by (11) we know that

$$\left| \Pr\{|H_n - pn| \geq \eta n\} \right| = \mathcal{O}\left(\exp(-c\eta^2 n)\right).$$

By the monotonicity of the distribution function this implies that for x with $|x - pn| \geq \eta n$ we also have

$$\left| \Pr\{|H_n - pn| \geq x\} \right| = \mathcal{O}\left(\exp(-c\eta^2 n)\right).$$

Thus (13) follows.

The first part of Theorem 1, i. e. (1), is now a trivial consequence of (12) and (13).

3. CONVERGENCE OF MOMENTS

Lemmata 2 and 3 can be easily used to get quite tight bounds for the expected value. Note that

$$\sum_{n \geq 0} \mathbb{E} H_n x^n = \sum_{k \geq 0} \left(\frac{1}{1-x} - y_k(x) \right). \quad (14)$$

Lemma 6. *The expected value of H_n can be bounded by*

$$pn + q \leq \mathbb{E} H_n \leq pn + \mathcal{O}\left(\frac{1}{p}\right). \quad (15)$$

Proof. Set

$$E(x) = \sum_{k \geq 0} \left(\frac{1}{1-x} - y_k(x) \right).$$

Then by (5) we obtain

$$E(x) \geq \frac{x}{1-x} \sum_{k \geq 0} \left(\frac{px}{1-qx} \right)^k = \frac{x(1-qx)}{(1-x)^2},$$

which is also true on the level of coefficients. Thus, we have

$$\mathbb{E} H_n \geq pn + q.$$

Similarly we get the upper bound. From

$$\begin{aligned} E(x) &\leq \frac{x(1-qx)}{(1-x)^2} + \frac{(qx)^2}{1-qx} \frac{1}{\left(\frac{px}{1-qx}\right) - qx} \left(\frac{1}{1 - \frac{px}{1-qx}} - \frac{1}{1-qx} \right) \\ &\quad + \frac{qx}{1-x} \left(\frac{px}{1-qx} \right) \frac{1}{\left(\frac{px}{1-qx}\right) - qx} \left(\frac{1}{1 - \frac{px}{1-qx}} - \frac{1}{1-qx} \right) \\ &\quad - \frac{qx^2}{1-x} \frac{1}{1-qx} \left(\frac{1}{1 - \frac{px}{1-qx}} - \frac{1}{1 - \frac{qx^2(1-q)}{1-qx}} \right) \\ &= \frac{x(1-qx)}{(1-x)^2} + \frac{(qx)^2}{(1-x)(1-qx)} \\ &\quad + \frac{pqx^2}{(1-x)^2(1-qx)} - \frac{qx^2}{(1-x)^2} + \frac{qx^2}{(1-x) \left(1 - \frac{pqx^2}{1-qx}\right)} \\ &= \frac{x(1-qx)}{(1-x)^2} + \frac{qx^2(1-qx)}{(1-x)(1-qx - pqx^2)} \end{aligned}$$

we directly obtain

$$\mathbb{E} H_n \leq pn + \mathcal{O}\left(\frac{1}{p}\right).$$

This completes the proof of Lemma 6. \square

Unfortunately it seems that we cannot prove similarly tight estimates for the variance. However, we can obtain a non-trivial result:

Lemma 7. *The variance of H_n can be bounded by*

$$\mathbb{V} H_n = pqn + \mathcal{O}_q\left(n^{1/2} \log^2 n\right). \quad (16)$$

Proof. If $F(x) = \mathbf{Pr}\{X \leq x\}$ denotes the distribution function of a random variable X (of compact support) then the variance of X can be represented by

$$\mathbb{V} X = 2 \int_{-\infty}^{\mathbb{E}X} (\mathbb{E}X - y)F(y) dy + 2 \int_{\mathbb{E}X}^{\infty} (y - \mathbb{E}X)(1 - F(y)) dy.$$

We apply this formula for the height H_n where we have very precise estimates for $\mathbb{E}H_n$ and its distribution function $F_n(x) = \mathbf{Pr}\{H_n \leq x\}$, compare with Lemma 5 and Lemma 6.

Let $u \leq \eta\sqrt{n}$ be a parameter to be defined later. By applying Lemma 5 and Lemma 6 we get

$$\begin{aligned} 2 \int_0^{\mathbb{E}H_n - u\sqrt{n}} (\mathbb{E}H_n - y)F_n(y) dy &= \mathcal{O}(ne^{-cu^2}), \\ 2 \int_{\mathbb{E}H_n - u\sqrt{n}}^{\mathbb{E}H_n} (\mathbb{E}H_n - y)F_n(y) dy &= \frac{pqn}{2} + \mathcal{O}(u^2\sqrt{n}), \\ 2 \int_{\mathbb{E}H_n}^{\mathbb{E}H_n - u\sqrt{n}} (y - \mathbb{E}H_n)(1 - F_n(y)) dy &= \frac{pqn}{2} + \mathcal{O}(u^2\sqrt{n}), \end{aligned}$$

and

$$2 \int_{\mathbb{E}H_n - u\sqrt{n}}^n (y - \mathbb{E}H_n)(1 - F_n(y)) dy = \mathcal{O}(ne^{-cu^2}).$$

Choosing $u = \log^2 n$ gives the result. \square

It should be further mentioned that it is quite easy to get asymptotic relations for all central moments of the form

$$\mathbb{E}(H_n - \mathbb{E}H_n)^k \sim c_k(pqn)^{k/2} \quad (n \rightarrow \infty)$$

by the method as described in Lemma 7, where $c_{2k} = (2k)!/(2^k k!)$ and $c_{2k+1} = 0$. Thus we have not only a weak convergence result for H_n (properly normalized) but convergence of moments, too.

REFERENCES

- [1] M. BARLOW, R. PEMANTLE AND E. PERKINS, *Diffusion limited aggregation on a homogeneous tree*, Prob. Theory and Related Fields **107** (1997), 1–60.
- [2] F. BERGERON, P. FLAJOLET AND B. SALVY, *Varieties of increasing trees*, Lecture Notes in Computer Science **581** (1992), 24–48.
- [3] L. DEVROYE AND B. REED, *On the variance of the height of random binary search trees*, SIAM J. Comput. **24** (1995), 1157–1162.
- [4] M. DRMOTA, *A bivariate asymptotic expansion of coefficients of powers of generating functions*, Europ. J. Combinatorics **15** (1994), 139–152.
- [5] M. DRMOTA, *The variance of the height of binary search trees*, Theoret. Comput. Sci. **270** (2002), 913–919.
- [6] B. REED, *The height of a random binary search tree*, J. Assoc. Comput. Mach., to appear.
- [7] W. FELLER, *An Introduction to Probability Theory and Its Applications, Vol. II*, 2nd ed., J. Wiley, New York, 1971.

- [8] H. PRODINGER, *A q -analogue of the path length of binary search trees*, *Algorithmica* **31** (2001), 433–441.

DEPARTMENT OF GEOMETRY, TECHNICAL UNIVERSITY OF VIENNA, WIEDNER HAUPTSTRASSE 8–10, A-1040 VIENNA, AUSTRIA

E-mail address: michael.drmota@tuwien.ac.at

THE JOHN KNOPFMACHER CENTRE FOR APPLICABLE ANALYSIS AND NUMBER THEORY, SCHOOL OF MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, P. O. WITS, 2050 JOHANNESBURG, SOUTH AFRICA

E-mail address: helmut@staff.ms.wits.ac.za

WWW-address: <http://www.wits.ac.za/helmut/index.htm>