

# Equivalents of NOTOP

Chris Laskowski  
University of Maryland

Joint work with Danielle Ulrich

Saharon's 80th Birthday Conference  
14 July, 2025

# History of Classification Theory



Chris Laskowski University of Maryland

Equivalents of NOTOP

# 1971 – Superstability

## Definition

A (complete, countable) stable theory  $T$  is **superstable** if there do not exist  $c$  and  $A_0 \subseteq A_1 \subseteq A_2 \subset \dots$  with  $\text{tp}(c/A_{n+1})$  forking over  $A_n$  for each  $n$ .

## Theorem

*If  $T$  is not superstable, then the class of uncountable models of  $T$  is chaotic. (In particular,  $I(T, \kappa) = 2^\kappa$  for all  $\kappa > \aleph_0$ .)*

**Henceforth**, we will assume all theories are (complete) and superstable in a countable language.

# T countable, superstable

## Notation:

M **a-model**  $\leftrightarrow \mathbf{F}_{\aleph_0}^a$ -saturated model  $\leftrightarrow \aleph_\epsilon$ -saturated model

means:  $M$  realizes every type in  $S(\text{acl}^{eq}(A))$  for every finite  $A \subseteq M$ .

An **independent triple of models**  $(M_0, M_1, M_2)$  satisfies  $M_0 \preceq M_1$ ,  $M_0 \preceq M_2$ , with  $M_1 \downarrow_{M_0} M_2$  An **independent triple of models**

$(M_0, M_1, M_2)$  satisfies  $M_0 \preceq M_1$ ,  $M_0 \preceq M_2$ , with  $M_1 \downarrow_{M_0} M_2$   
(**forking independence!**)

# 1981 – NDOP

## Definition

A (countable) superstable  $T$  has **NDOP** if, for any independent triple  $(M_0, M_1, M_2)$  of  $a$ -models, any  $a$ -prime model  $M^*$  over  $M_1 M_2$  is minimal over  $M_1 M_2$ . [If  $M_1 M_2 \subseteq N \preceq M^*$ , then  $N = M^*$ .]

## Theorem (Main Gap for $a$ -saturated models)

*If  $T$  is superstable with NDOP, then every  $a$ -model is  $a$ -prime and  $a$ -minimal over an independent tree  $\{M_\eta : \eta \in I\}$  of  $a$ -models of size  $2^{\aleph_0}$ .*

## Theorem

*If  $T$  is either unsuperstable or if  $T$  has DOP, then  $I(T, \kappa) = 2^\kappa$  for all  $\kappa > \aleph_0$ .*

# 1984 – The ‘Magic Bullet’

## Definition

A (countable) superstable  $T$  has **NOTOP** if there **does not** exist a type  $p(x, y, z)$  such that for every  $\lambda$  and  $R \subseteq \lambda^2$ , there is a model  $M_R$  and  $\{a_i : i \in \lambda\} \subseteq M_R$  such that for all  $(i, j) \in \lambda^2$ ,

$M_R$  realizes  $p(x, a_i, a_j)$  if and only if  $R(i, j)$

## Definition

$T$  is **classifiable** if  $T$  is countable, superstable, NDOP, NOTOP.

## 1989 – 2 years after Volume 2 of Classification Theory

### Theorem

*Let  $T$  be any complete theory in a countable language.*

- 1 *If  $T$  is not classifiable, the  $I(T, \kappa) = 2^\kappa$  for all  $\kappa > \aleph_0$ .*
- 2 *if  $T$  is classifiable, then every model  $N$  is **constructible** and **minimal** over an independent tree  $(M_\eta : \eta \in I)$  of countable, elementary substructures.*

Computing the 13 species of uncountable spectra starts with this.

# The take-away

Historically –

NOTOP was only developed/explored in the presence of NDOP!

**Will see:** Countable, superstable, NOTOP theories admit structure theorems, even without NDOP.



# Notions of isolation

**Recall:**  $\text{tp}(c/B)$  is **isolated** if there is some  $\psi(x, b) \in \text{tp}(c/B)$  such that  $\psi(x, b) \vdash \text{tp}(c/B)$ .

A weakening:

**Lachlan:**  $\text{tp}(c/B)$  is  **$\ell$ -isolated** if, for every  $\phi(x, y)$ , there is  $\psi(x, b) \in \text{tp}(c/B)$  such that  $\psi(x, b) \vdash \text{tp}_\phi(c/B)$ .

- If  $T$  is  $\omega$ -stable, then for every set  $B$ , the isolated types are dense in  $S(B)$ .
- If  $T$  is countable, superstable, then for every set  $B$ , the  $\ell$ -isolated types are dense in  $S(B)$ .

## Theorem (L-Ulrich)

For  $T$  countable, superstable, TFAE:

- 1 NOTOP;
- 2 For all independent triples  $(M_0, M_1, M_2)$  and all finite  $c$ ,  
 $\text{tp}(c/M_1M_2)$   *$\ell$ -isolated implies*  $\text{tp}(c/M_1M_2)$  *isolated*.

**Recall:**  $T$  is classifiable iff every  $N \models T$  is **constructible** and **minimal** over an independent tree  $(M_\eta : \eta \in I)$  of countable, elementary substructures.

### Theorem (L-Ulrich)

*Suppose  $T$  is countable, superstable, with NOTOP.*

- 1 Every  $N \models T$  is **atomic** over an independent tree  $(M_\eta : \eta \in I)$  of countable, elementary substructures;
- 2 There is a constructible model  $N_0 \preceq N$  over  $\bigcup\{M_\eta : \eta \in I\}$ ;
- 3 If  $N_0 \preceq N_1 \preceq N$ , then  $N_0 \preceq_{\infty, \omega} N_1 \preceq_{\infty, \omega} N$ , i.e., all three models are back-and-forth equivalent.

## Contrast:

- If  $T$  is classifiable, then every model  $N$  has a tree  $(M_\eta : \eta \in I)$  of countable, elementary substructures that determines  $N$  up to isomorphism over the tree.
- If  $T$  is countable, superstable, NOTOP, then every model  $N$  has a tree  $(M_\eta : \eta \in I)$  of countable, elementary substructures that determines  $N$  up to back and forth equivalence over the tree.

## 'Under the hood' – Study independent triples of models

Say  $\overline{M} = (M_0, M_1, M_2)$ ,  $\overline{N} = (N_0, N_1, N_2)$  are independent triples of models (of any size). Define  $\overline{M} \sqsubseteq \overline{N}$  iff  $M_i \preceq N_i$  for each  $i$ ,  $N_0 \downarrow_{M_0} M_1 M_2$ ,  $N_1 \downarrow_{N_0 M_1} M_2$  and  $N_2 \downarrow_{N_0 M_2} M_1$ .

**Credo:** (Indep triples,  $\sqsubseteq$ ) acts very much like  $(\text{Mod}(\mathcal{T}), \preceq)$ .

- If  $\overline{M} \sqsubseteq \overline{N}$  then  $M_1 M_2 \subseteq_{TV} N_1 N_2$ ;
- (ULS) For any  $\overline{M}$ , there is  $\overline{N} \sqsupseteq \overline{M}$  consisting of a-models
- (DLS) For any  $\overline{N}$  and any  $X \subseteq N_1 N_2$  with  $|X| \leq \kappa$ , there is  $\overline{M} \sqsubseteq \overline{N}$  with  $X \subseteq M_1 M_2$  and  $|M_1 M_2| \leq \kappa$ .

### Definition (Harrington)

Suppose  $\overline{M} = (M_0, M_1, M_2)$  is any independent triple. We say  $c$  is  $V$ -dominated by  $\overline{M}$  if,  $c \perp_{M_1 M_2} N_1 N_2$  for every  $\overline{N} \supseteq \overline{M}$ .

**New:** We say  $T$  has  $V$ -DI if for all  $c$  and for all  $\overline{M}$ , if  $c$  is  $V$ -dominated by  $\overline{M}$ , then  $\text{tp}(c/M_1 M_2)$  is isolated.

**Fact:** For any  $c$  and  $\overline{M}$ ,

- If  $\text{tp}(c/M_1 M_2)$  is  $\ell$ -isolated, then  $c$  is  $V$ -dominated by  $\overline{M}$ .
- If, in addition, each  $M_i$  is  $\mathbf{F}_{\aleph_0}^a$ -saturated, then the converse holds.

**Will see:**  $V$ -DI is another equivalent of NOTOP.

# Two consequences of V-DI

One consequence:

Theorem (L-Ulrich)

*V-DI implies PMOP (existence of a constructible model over independent triples of models of any size).*

**Remark:** The above was proved by Shelah, and reproved by Hart, both under the assumption of NDOP.

On page 619 of *Classification Theory* (1987), Shelah writes:

“Remark. Really “without the dop” is not necessary, this will be shown in a subsequent paper.”

# Local versions of NDOP

**Fact:**  $T$  has NDOP iff for all independent triples  $(M_0, M_1, M_2)$  of  $\mathcal{A}$ -models and for all  $\mathcal{A}$ -prime  $M^*$  over  $M_1 M_2$ , every regular type  $r \not\perp M^*$  is  $\not\perp M_1$  or  $\not\perp M_2$ .

**Fact:**  $\not\perp$  induces an equivalence relation on the set of regular types.

Let  $\mathbf{P}$  be any union of  $\not\perp$ -classes of regular types.

**Definition:**  $T$  has  $\mathbf{P}$ -NDOP iff for all independent triples  $(M_0, M_1, M_2)$  of  $\mathcal{A}$ -models and for all  $\mathcal{A}$ -prime  $M^*$  over  $M_1 M_2$ , every regular  $r \not\perp M^*$  with  $r \in \mathbf{P}$  is  $\not\perp M_1$  or  $\not\perp M_2$ .

[L-Shelah] For sufficiently nice  $\mathbf{P}$ ,  $\mathcal{A}$ -models of superstable  $T$  with  $\mathbf{P}$ -NDOP admit decomposition trees.



# A new class of regular types

**Definition** (Baisalov, 1990) An **e-type** is a stationary, weight one type  $p(x, d)$  with  $d$  finite that is non-isolated.

**Definition**  $\mathbf{P}_e = \{\text{regular } r : r \not\vdash \text{some } e\text{-type } p(x, d)\}$ .

**Note:**  $\mathbf{P}_e$ -NDOP is a slight strengthening of eni-NDOP (they are equivalent if  $T$  is  $\omega$ -stable).

**Second consequence:**

- For  $T$  countable, superstable, V-DI implies  $\mathbf{P}_e$ -NDOP.

# Some equivalents

## Theorem (L-Ulrich)

*The following are equivalent for a countable, superstable  $T$ :*

- 1 For every independent triple of *countable* models  $\overline{M}$ ,  $\text{tp}(c/M_1M_2)$   $\ell$ -isolated implies  $\text{tp}(c/M_1M_2)$  isolated;
- 2  $T$  is V-DI;
- 3  $T$  has  $\mathbf{P}_e$ -NDOP and countable PMOP (there exists a constructible model over every independent triple of *countable* models);
- 4  $T$  has  $\mathbf{P}_e$ -NDOP and full PMOP (there exists a constructible model over every independent triple of models);
- 5  $T$  has NOTOP.

## On adding constants

**Recall:**  $T$  has OTOP iff if there is a type  $p(x, y, z)$  such that for every  $\lambda$  and  $R \subseteq \lambda^2$ , there is a model  $M_R$  and  $\{a_i : i \in \lambda\} \subseteq M_R$  such that for all  $(i, j) \in \lambda^2$ ,

$$M_R \text{ realizes } p(x, a_i, a_j) \text{ if and only if } R(i, j)$$

**Good news:** If a countable, superstable theory has OTOP, then any expansion by adding **countably many** constants will also have OTOP. (hence, we may assume our type  $p(x, y, z)$  witnessing OTOP has countably many parameters).

**Danger:** There is a countable, superstable theory  $T$  with OTOP, but if we add  $2^{\aleph_0}$  constants naming a saturated model, then the expanded theory is categorical in all  $\kappa > 2^{\aleph_0}$ .

## On a personal note

Thank you Saharon,  
for all the time and energy you spent mentoring me.

And thanks to all of you for listening!