

ON WELL-SPLITTING POSETS

DUŠAN REPOVŠ AND LYUBOMYR ZDOMSKYY

ABSTRACT. We introduce a class of proper posets which is preserved by countable support iterations, includes ω^ω -bounding, Cohen, Miller, and Mathias posets associated to filters with the Hurewicz covering properties, and has the property that the ground model reals remain splitting and unbounded in corresponding extensions.

1. INTRODUCTION

The famous Roitman problem asks whether it is consistent, relative to the consistency of ZFC, that $\mathfrak{d} = \omega_1 < \mathfrak{a}$. Here, \mathfrak{d} is the minimal cardinality of a subfamily of ω^ω which is *dominating* with respect to the preorder relation \leq^* on ω^ω , where $a \leq^* b$ for $a, b \in \omega^\omega$ means that $a(n) \leq b(n)$ for all but finitely many n ; \mathfrak{a} is the minimal cardinality of a *mad subfamily* \mathcal{A} of $[\omega]^\omega$, i.e., a subfamily whose distinct elements have finite intersection and which is maximal with respect to this property.

Without the restriction $\mathfrak{d} = \omega_1$, the consistency of $\mathfrak{d} < \mathfrak{a}$ has been established in a breakthrough work of Shelah [12]. Regarding the original Roitman problem, even the following weaker version raised in [4] remains open: Is it consistent that $\mathfrak{s} = \mathfrak{b} = \omega_1 < \mathfrak{a}$? Here, \mathfrak{s} is the minimal cardinality of a *splitting* family, i.e., a family $\mathcal{S} \subset [\omega]^\omega$ such that for every $X \in [\omega]^\omega$ there exists $S \in \mathcal{S}$ for which both $S \cap X$ and $X \setminus S$ are infinite; \mathfrak{b} is the minimal cardinality of a subfamily of ω^ω which is *unbounded* with respect to \leq^* . It is well-known that $\max\{\mathfrak{b}, \mathfrak{s}\} \leq \mathfrak{d}$ and the strict inequality holds, e.g., in the Cohen model (see [2, 13] for more information on these and many other cardinal characteristics of the continuum).

In this note we isolate the class of well-splitting posets (see the next section for the definition) with properties described in the abstract, aiming at the solution of the aforementioned weak version of Roitman's problem. This class includes among others Mathias posets associated to filters on ω with the Hurewicz covering property. This motivates the following

Question 1.1. (CH) Can every mad family be destroyed by a well-splitting poset? In particular, given a mad family \mathcal{A} , is there a well-splitting poset \mathbb{P} such that in $V^{\mathbb{P}}$, $\{\omega \setminus A : A \in \mathcal{A}\}$ can be enlarged to a Hurewicz filter, or more generally to a filter, whose Mathias forcing is well-splitting?

2010 *Mathematics Subject Classification.* Primary: 03E35, 03E17; Secondary: 54D20.

Key words and phrases. Splitting, bounding, Miller forcing, filter, Hurewicz space, mad family.

The first author was partially supported by the Slovenian Research Agency grants P1-0292, N1-0114, and N1-0083. The second author would like to thank the Austrian Science Fund FWF (Grants I2374-N35 and I3709-N35) for generous support for this research.

By Theorem 2.7 proved in the next section, the affirmative answer to Question 1.1 would allow to construct a model of $\mathfrak{b} = \mathfrak{s} = \omega_1 < \mathfrak{a} = \omega_2$.

Recall from [8] that a topological space X is said to have the *Hurewicz covering property* (or is simply called Hurewicz) if for every sequence $\langle \mathcal{U}_n : n \in \omega \rangle$ of open covers of X there exists a sequence $\langle \mathcal{V}_n : n \in \omega \rangle$ such that each \mathcal{V}_n is a finite subfamily of \mathcal{U}_n and the collection $\{\cup \mathcal{V}_n : n \in \omega\}$ is a γ -cover of X , i.e., the set $\{n \in \omega : x \notin \cup \mathcal{V}_n\}$ is finite for each $x \in X$. It is clear that σ -compact spaces are Hurewicz, but by [9, Theorem 5.1] there exist also non- σ -compact sets of reals having the Hurewicz property. We consider each filter on ω with the subspace topology inherited from $\mathcal{P}(\omega)$, the latter being a topological copy of the Cantor space 2^ω via characteristic functions. As it was proved in [7], the Mathias forcing associated to a filter \mathcal{F} is almost ω^ω -bounding in terminology of [11] if and only if \mathcal{F} is Hurewicz. It is worth mentioning here that in general, almost ω^ω -bounding posets can make ground model reals non-splitting, see, e.g., [11, Lemma 1.14], so by Lemma 2.1 almost ω^ω -bounding posets do not have to be well-splitting.

Built on the proof of [3, Theorem 3.1], it is established in [14] that under CH, for every mad family \mathcal{A} , the collection $\{\omega \setminus A : A \in \mathcal{A}\}$ can be enlarged to an ultrafilter \mathcal{F} with a certain covering property which is weaker (but similar) to the Hurewicz one, and whose Mathias forcing does not produce any new real dominating the given ground model unbounded subset. The construction in [14] cannot be directly used to answer Question 1.1 since by Lemma 2.1 the Mathias forcing for ultrafilters cannot be well-splitting because it adds an unsplit real. However, it is natural to ask how far can we weaken the Hurewicz property of a filter so that its Mathias forcing is still well-splitting.

Question 1.2. Let \mathcal{F} be a filter on ω whose Mathias forcing is well-splitting. Is then \mathcal{F} Hurewicz? In other words, are being well-splitting and almost ω^ω -bounding equivalent for such posets?

2. WELL-SPLITTING POSETS

Throughout this section we denote by E_0 and E_1 the sets of all even and odd natural numbers, respectively. A strictly increasing function $f \in \omega^\omega$ is said to *well-split* a set M if the sets $\{n \in E_j : |[f(n), f(n+1)) \cap Y| \geq 2\}$ are infinite for all $Y \in M \cap [\omega]^\omega$ and $j \in 2$.

We shall say that a poset \mathbb{P} is *well-splitting* if the following is satisfied: Whenever $\mathbb{P} \in M$, where M is a countable elementary submodel of $H(\theta)$ for any sufficiently large θ , $p \in M \cap \mathbb{P}$ and f well-splits M , then there is some $q \leq p$ which is (M, \mathbb{P}) -generic and such that q forces f to well-split $M[\Gamma]$, where Γ is the canonical name for \mathbb{P} -generic filter.

Lemma 2.1. *Suppose that \mathbb{P} is well-splitting and G is \mathbb{P} -generic. Then $V \cap [\omega]^\omega$ is splitting and $V \cap \omega^\omega$ is unbounded in $V[G]$.*

Proof. To see that $\omega^\omega \cap V$ is unbounded, let us fix a \mathbb{P} -name \dot{h} for an element of $\omega^{\uparrow\omega}$ (the family of all strictly increasing functions in ω^ω), a countable elementary submodel M of $H(\theta)$ such that $\mathbb{P}, \dot{h} \in M$, and $p \in \mathbb{P} \cap M$.

Suppose that f well-splits M and $q \leq p$ is any (M, \mathbb{P}) -generic condition which forces f to well-split $M[\Gamma]$. Let $\dot{h}_1 \in M$ be a \mathbb{P} -name for the following function: $\dot{h}_1(0) = 0$, $\dot{h}_1(n+1) = \dot{h}_1(\dot{h}_1(n)) + 1$ for all $n \in \omega$. It follows from the above that q forces the set

$$\dot{I} := \{n \in E_0 : |[f(n), f(n+1)) \cap \text{range}(\dot{h}_1)| \geq 2\}$$

to be infinite. Let $G \ni q$ be \mathbb{P} -generic and set $I = \dot{I}^G$, $h = \dot{h}_1^G$, and $h_1 = \dot{h}_1^G$. In $V[G]$, for every $i \in I$ we can find $n_i \in \omega$ such that $h_1(n_i), h_1(n_i+1) \in [f(i), f(i+1))$. Thus $h_1(n_i+1) = h(h_1(n_i)) < f(i+1) \leq f(h_1(n_i))$, i.e.,

$$\{h_1(n_i) : i \in \omega\} \subset \{k : h(k) < f(k)\},$$

and hence h does not dominate f . Summarizing the above, we conclude that for any $p \in \mathbb{P}$ and any \mathbb{P} -name \dot{h} for an element of $\omega^{\uparrow\omega}$, there is a stronger condition q and $f \in \omega^{\uparrow\omega} \cap V$ such that q forces the set $\{k : \dot{h}(k) < f(k)\}$ to be infinite. This precisely means that $\omega^{\uparrow\omega} \cap V$ is unbounded in $V[G]$ for any \mathbb{P} -generic filter G .

To prove that $[\omega]^\omega \cap V$ is splitting, let us fix a \mathbb{P} -name \dot{Y} for an element of $[\omega]^\omega$, a countable elementary submodel M of $H(\theta)$ such that $\mathbb{P}, \dot{Y} \in M$, and $p \in \mathbb{P} \cap M$. Suppose that f well-splits M and $q \leq p$ is any (M, \mathbb{P}) -generic condition which forces f to well-split $M[\Gamma]$. Then q forces the sets

$$\dot{I}_j := \bigcup_{n \in E_j} [f(n), f(n+1)) \cap \dot{Y}$$

to be infinite for all $j \in 2$. Since the sets $\bigcup_{n \in E_j} [f(n), f(n+1))$, $j \in 2$, are disjoint, infinite, and both are in V , this completes our proof. \square

It is clear that each well-splitting poset is proper and an iteration of finitely many well-splitting posets is again well-splitting. Next, we shall establish that being well-splitting is also preserved by countable support iterations. The proof of the following lemma is similar to that of [1, Lemma 2.8], with some additional control on the sequence $\langle \dot{p}_i : i \in \omega \rangle$.

Let us make a couple of standard conventions regarding our notation. Whenever $\langle \mathbb{P}_\alpha, \dot{Q}_\alpha : \alpha < \delta \rangle$ is an iterated forcing construction, we denote by $\mathbb{P}_{[\alpha_0, \alpha_1]}$ a \mathbb{P}_{α_0} -name for the quotient poset $\mathbb{P}_{\alpha_1}/\mathbb{P}_{\alpha_0}$, viewed naturally as an iteration over the ordinals $\xi \in \alpha_1 \setminus \alpha_0$. For a \mathbb{P}_{α_0} -generic G and a \mathbb{P}_{α_1} -name τ , where $\alpha_0 \leq \alpha_1$, we denote by τ^G the $\mathbb{P}_{[\alpha_0, \alpha_1]}^G$ -name in $V[G]$ obtained from τ by partially interpreting it with G . This allows us to speak about, e.g., $\mathbb{P}_{[\alpha_1, \alpha_2]}^G$ for $\alpha_0 \leq \alpha_1 \leq \alpha_2 \leq \delta$ and a \mathbb{P}_{α_0} -generic filter G . For a poset \mathbb{P} we shall denote by $\Gamma_{\mathbb{P}}$ the standard \mathbb{P} -name for \mathbb{P} -generic filter. We shall write Γ_α instead of $\Gamma_{\mathbb{P}_\alpha}$ whenever we work with an iterated forcing construction which is clear from the context. Also, $\Gamma_{[\alpha_0, \alpha_1]}$ is a \mathbb{P}_{α_1} -name whose interpretation with respect to a \mathbb{P}_{α_0} -generic filter G is $\Gamma_{\mathbb{P}_{[\alpha_0, \alpha_1]}^G}$, which is an element of $V[G]$.

Lemma 2.2. *If $\langle \mathbb{P}_\alpha, \dot{Q}_\alpha : \alpha < \delta \rangle \in M$ is a countable support iteration of well-splitting (hence proper) posets, then \mathbb{P} is also well-splitting.*

Proof. The proof is by induction on δ . The successor case is clear. So assume that δ is limit, $p \in \mathbb{P}_\delta$, and $M \ni \mathbb{P}_{\delta,p}$ is a countable elementary submodel of $H(\theta)$ for a sufficiently large θ . Pick an increasing sequence $\langle \delta_i : i \in \omega \rangle$ cofinal in $\delta \cap M$, with $\delta_i \in M$ for all $i \in \omega$. Let also $\{D_i : i \in \omega\}$ and $\{\dot{Y}_i : i \in \omega\}$ be an enumeration of all open dense subsets of \mathbb{P}_δ and all \mathbb{P}_δ -names for an infinite subset of ω which are elements of M , respectively. We can assume without loss of generality, that for every \mathbb{P}_δ -name $\dot{Y} \in M$ for an element of $[\omega]^\omega$ the set $\{i \in \omega : \dot{Y} = \dot{Y}_i\}$ is infinite. Suppose that f well-splits M . We will define by induction on $i \in \omega$ a condition $q_i \in \mathbb{P}_{\delta_i}$ and \mathbb{P}_{δ_i} -name $\dot{p}_i, \dot{n}_i^0, \dot{n}_i^1$ such that

- (i) \dot{p}_i is a name for an element of \mathbb{P}_δ , $q_0 \Vdash_{\delta_0} \dot{p}_0 \leq \check{p}$, and $q_{i+1} \Vdash_{\delta_{i+1}} \dot{p}_{i+1} \leq \dot{p}_i$;
- (ii) $q_{i+1} \upharpoonright \delta_i = q_i$;
- (iii) q_i is $(M, \mathbb{P}_{\delta_i})$ -generic;
- (iv) \dot{n}_i^0, \dot{n}_i^1 are \mathbb{P}_{δ_i} -names for natural numbers bigger than i ; and
- (v) q_i forces over \mathbb{P}_{δ_i} that “ $\dot{p}_i \upharpoonright \delta_i \in \Gamma_{\delta_i}$, $\dot{p}_i \in D_i \cap M$, and \dot{p}_i forces over \mathbb{P}_δ that $\dot{n}_i^j \in E_j$ and $|\{f(\dot{n}_i^j), f(\dot{n}_i^j + 1)\} \cap \dot{Y}_i| \geq 2$ for all $j \in 2$ ”.

Suppose now that we have constructed objects as above and set $q = \bigcup_{i \in \omega} q_i$. Since $q_i = q \upharpoonright \delta_i$ forces over \mathbb{P}_{δ_i} that $\dot{p}_i \upharpoonright \delta_i \in \Gamma_{\delta_i}$ and $q_{i+1} \Vdash_{\delta_{i+1}} \dot{p}_{i+1} \leq \dot{p}_i$ for all i , a standard argument yields that q is (M, \mathbb{P}_δ) -generic and $q \Vdash_\delta \dot{p}_i \in \Gamma_\delta$ for all $i \in \omega$, see, e.g., the proof of [1, Lemma 2.8] for details. Then q forces that $\tau_0 := \{n \in E_0 : |\{f(n), f(n+1)\} \cap \dot{Y}| \geq 2\}$ and $\tau_1 := \{n \in E_1 : |\{f(n), f(n+1)\} \cap \dot{Y}| \geq 2\}$ are infinite for any \mathbb{P}_δ -name \dot{Y} for an infinite subset of ω : Given \mathbb{P}_δ -generic $G \ni q$, note that $p_i := \dot{p}_i^G \in G$ for all i . Now (v) implies $n_i^j \in \tau_j^G$ for all $j \in 2$ and $i \in \omega$ such that $\dot{Y} = \dot{Y}_i$, where $n_i^j = (\dot{n}_i^j)^G$.

Returning now to the inductive construction, assume that $q_i \in \mathbb{P}_{\delta_i}$, \mathbb{P}_{δ_i} -names $\dot{p}_i, \dot{n}_i^0, \dot{n}_i^1$ satisfying (i)-(v) have already been constructed. Let G_{δ_i} be \mathbb{P}_{δ_i} -generic containing q_i and $p_i = \dot{p}_i^{G_{\delta_i}} \in \mathbb{P}_\delta \cap M$. By (v) we know that $p_i \upharpoonright \delta_i \in G_{\delta_i}$. In $V[G_{\delta_i}]$ let $p'_i \in M \cap D_{i+1}$ be such that $p'_i \leq p_i$ and $p'_i \upharpoonright \delta_i \in G_{\delta_i}$. By the maximality principle we get a \mathbb{P}_{δ_i} -name \dot{p}'_i for a condition in \mathbb{P}_δ such that $q_i \Vdash_{\delta_i}$ “ $\dot{p}'_i \leq \dot{p}_i$, $\dot{p}'_i \in M \cap D_{i+1}$, and $\dot{p}'_i \upharpoonright \delta_i \in \Gamma_{\delta_i}$ ”.

Given a $\mathbb{P}_{\delta_{i+1}}$ -generic filter R , in $V[R]$ construct a decreasing sequence $\langle r_m : m \in \omega \rangle \in M[R]$ of conditions in $\mathbb{P}_{[\delta_{i+1}, \delta]}^R$ below $(\dot{p}'_i \upharpoonright [\delta_{i+1}, \delta])^R$ such that for some $a_m \in [\omega]^m$ we have $r_m \Vdash_{\mathbb{P}_{[\delta_{i+1}, \delta]}^R}$ “ a_m is the set of the first m elements of \dot{Y}_{i+1} ”. By the maximality principle we get a sequence $\langle \rho_m : m \in \omega \rangle \in M$ of $\mathbb{P}_{\delta_{i+1}}$ -names for elements of $\mathbb{P}_{[\delta_{i+1}, \delta]}$ such that

$$\Vdash_{\delta_{i+1}} [\rho_{m+1} \leq \rho_m \wedge \exists \nu_m \in [\omega]^m (\rho_m \Vdash_{\mathbb{P}_{[\delta_{i+1}, \delta]}} \nu_m \text{ is the set of the first } m \text{ many elements of } \dot{Y}_{i+1})].$$

In the notation used above, let \dot{Z} be a $\mathbb{P}_{\delta_{i+1}}$ -name for $\bigcup_{m \in \omega} \nu_m$ and note that \dot{Z} is a $\mathbb{P}_{\delta_{i+1}}$ -name for an infinite subset of ω .

Let again G_{δ_i} be \mathbb{P}_{δ_i} -generic containing q_i and $p'_i = (\dot{p}'_i)^{G_{\delta_i}} \in \mathbb{P}_\delta \cap M \cap D_{i+1}$. It also follows from the above that $p'_i \upharpoonright \delta_i \in G_{\delta_i}$. For a while we shall

work in $V[G_{\delta_i}]$. Since $\mathbb{P}_{[\delta_i, \delta_{i+1}]}^{G_{\delta_i}}$ is well-splitting in $V[G_{\delta_i}]$ by our inductive assumption, there exists a $(M[G_{\delta_i}], \mathbb{P}_{[\delta_i, \delta_{i+1}]}^{G_{\delta_i}})$ -generic condition $\pi \leq p'_i \upharpoonright [\delta_i, \delta_{i+1}]^{G_{\delta_i}}$ such that

$$\pi \Vdash_{\mathbb{P}_{[\delta_i, \delta_{i+1}]}^{G_{\delta_i}}} \tau_j := \{n \in E_j : |[f(n), f(n+1)] \cap \dot{Z}^{G_{\delta_i}}| \geq 2\}$$

is infinite for all $j \in 2$. Let H be $\mathbb{P}_{[\delta_i, \delta_{i+1}]}^{G_{\delta_i}}$ -generic over $V[G_{\delta_i}]$ containing π and $n_{i+1}^j \in \tau_j^H \setminus (i+2)$, where $j \in 2$. In $V[G_{\delta_i} * H]$ pick $m \in \omega$ such that

$$r_m := \rho_m^{G_{\delta_i} * H} \Vdash_{\mathbb{P}_{[\delta_{i+1}, \delta]}^{G_{\delta_i} * H}} \dot{Z}^{G_{\delta_i} * H} \cap f(\max_{j \in 2}(n_{i+1}^j) + 1) = \dot{Y}_{i+1}^{G_{\delta_i} * H} \cap f(\max_{j \in 2}(n_{i+1}^j) + 1).$$

In $M[G_{\delta_i}]$ pick a condition $s \in M[G] \cap H$ below $p'_i \upharpoonright [\delta_i, \delta_{i+1}]^{G_{\delta_i}}$ forcing the above properties of n_{i+1}^j , τ_j , and ρ_m , where $j \in 2$. By the maximality principle we obtain $\mathbb{P}_{[\delta_i, \delta_{i+1}]}^{G_{\delta_i}}$ -names \dot{s} and ρ in $M[G_{\delta_i}]$ for element of $\mathbb{P}_{[\delta_i, \delta_{i+1}]}^{G_{\delta_i}}$ and $\mathbb{P}_{[\delta_{i+1}, \delta]}^{G_{\delta_i}}$, and names \dot{n}_{i+1}^j for natural numbers such that

$$(1) \quad \begin{aligned} \pi \Vdash_{\mathbb{P}_{[\delta_i, \delta_{i+1}]}^{G_{\delta_i}}} \dot{s} \in M[G_{\delta_i}] \cap \Gamma_{[\delta_i, \delta_{i+1}]}^{G_{\delta_i}} \wedge \dot{s} \leq \dot{p}' \upharpoonright [\delta_i, \delta_{i+1}]^{G_{\delta_i}} \wedge \dot{s} \Vdash_{\mathbb{P}_{[\delta_i, \delta_{i+1}]}^{G_{\delta_i}}} \\ \rho \leq \dot{p}' \upharpoonright [\delta_{i+1}, \delta]^{G_{\delta_i}} \wedge \rho \Vdash_{\mathbb{P}_{[\delta_{i+1}, \delta]}^{G_{\delta_i}}} \\ \forall j \in 2 \ |[f(\dot{n}_{i+1}^j), f(\dot{n}_{i+1}^j + 1)] \cap \dot{Y}_{i+1}| \geq 2. \end{aligned}$$

Using the maximality principle again, we can find \mathbb{P}_{δ_i} -names for the objects appearing in equation (1) such that q_i forces this equation. We shall use the same notation for these names. It remains to set $q_{i+1} = q_i \hat{\wedge} \pi$ and $\dot{p}_{i+1} = \dot{p}' \upharpoonright \delta_i \hat{\wedge} \dot{s} \hat{\wedge} \rho$ and note that they together with the names \dot{n}_{i+1}^j , $j \in 2$, satisfy (i)-(v) for $i+1$. \square

By a Miller tree we understand a subtree T of $\omega^{<\omega}$ consisting of increasing finite sequences such that the following conditions are satisfied:

- Every $t \in T$ has an extension $s \in T$ which is splitting in T , i.e., there are more than one immediate successors of s in T ;
- If s is splitting in T , then it has infinitely many successors in T .

The Miller forcing is the collection \mathbb{M} of all Miller trees ordered by inclusion, i.e., smaller trees carry more information about the generic. This poset was introduced in [10]. For a Miller tree T we shall denote by $\text{Split}(T)$ the set of all splitting nodes of T . $\text{Split}(T)$ may be written in the form $\bigcup_{i \in \omega} \text{Split}_i(T)$, where

$$\text{Split}_i(T) = \{t \in \text{Split}(T) : |\{s \in \text{Split}(T) : s \subsetneq t\}| = i\}.$$

If $T_0, T_1 \in \mathbb{M}$, then $T_1 \leq_i T_0$ means $T_1 \leq T_0$ and $\text{Split}_i(T_1) = \text{Split}_i(T_0)$. It is easy to check that for any sequence $\langle T_i : i \in \omega \rangle \in \mathbb{M}^\omega$, if $T_{i+1} \leq_i T_i$ for all i , then $\bigcap_{i \in \omega} T_i \in \mathbb{M}$.

For a node t in a Miller tree T we denote by T_t the set $\{s \in T : s \text{ is compatible with } t\}$. It is clear that T_t is also a Miller tree.

Lemma 2.3. *The Miller forcing \mathbb{M} is well-splitting.*

Proof. Let N be an elementary submodel of $H(\theta)$ and $T \in \mathbb{M} \cap N$. Let $\{\dot{Y}_i : i \in \omega\}$ be an enumeration of all \mathbb{M} -names for infinite subsets of ω which are elements of N , in which every such name appears infinitely often. Let also $\{D_i : i \in \omega\}$ be an enumeration of all open dense subsets of \mathbb{M} which belong to N . Suppose that $f \in \omega^\omega$ well-splits N . We shall inductively construct a sequence $\langle T_i : i \in \omega \rangle$ such that $T_{i+1} \leq_i T_i$ and $T_\infty = \bigcap_{i \in \omega} T_i$ is as required. Set $T_0 = T$ and suppose that T_i has already been constructed. Moreover, we shall assume that $(T_i)_t \in N$ for all $t \in \text{Split}_i(T_i)$. Let $\{t_j : j \in \omega\}$ be a bijective enumeration of $\text{Split}_i(T_i)$. For every j and $k \in \omega$ such that $t_j \hat{\ } k \in T_i$ fix a decreasing sequence $\langle S_n^{i,j,k} : n \in \omega \rangle \in N$ of elements of D_i below $(T_i)_{t_j \hat{\ } k}$ such that each $S_n^{i,j,k}$ decides some $a_n^{i,j,k} \in [\omega]^n$ to be the set of the first n many elements of \dot{Y}_i . Thus $Y^{i,j,k} := \bigcup_{n \in \omega} a_n^{i,j,k} \in N \cap [\omega]^\omega$, and hence there are $E_p \ni m_{n,p}^{i,j,k} \geq i$ such that

$$|[f(m_{n,p}^{i,j,k}), f(m_{n,p}^{i,j,k} + 1)) \cap Y_n^{i,j,k}] \geq 2$$

for all $p \in 2$. Let $n(i, j, k)$ be such that

$$Y^{i,j,k} \cap \max_{p \in 2} f(m_{n(i,j,k),p}^{i,j,k} + 1) \subset a_{n(i,j,k)}^{i,j,k}$$

and set

$$T_{i+1} = \bigcup \{S_{n(i,j,k)}^{i,j,k} : j \in \omega, t_j \hat{\ } k \in T_i\}.$$

This completes our inductive construction of the fusion sequence $\langle T_i : i \in \omega \rangle$. We claim that T_∞ is as required. First of all, T_∞ is (N, \mathbb{M}) -generic because the collection $\bigcup \{S_{n(i,j,k)}^{i,j,k} : j \in \omega, t_j \hat{\ } k \in T_i\}$ is a subset of D_i and predense below T_{i+1} (and hence also below T_∞). Now fix a \mathbb{M} -name $\dot{Y} \in N$ for an element of $[\omega]^\omega$ and suppose to the contrary, that there exist $i \in \omega$, $p \in 2$, and $R \leq T_\infty$ that forces $|[f(m), f(m+1)) \cap \dot{Y}] \leq 1$ for all $E_p \ni m \geq i$. Enlarging i , if necessary, we may assume that $\dot{Y} = \dot{Y}_i$. Passing to a stronger condition, if necessary, we may assume that $R \leq (T_i)_{t_j \hat{\ } k}$ for some $i, j \in \omega$ and k such that $t_j \hat{\ } k \in T_i$. But then $R \leq S_{n(i,j,k)}^{i,j,k}$, and the latter condition forces

$$|[f(m_{n,p}^{i,j,k}), f(m_{n,p}^{i,j,k} + 1)) \cap Y_n^{i,j,k}] = |[f(m_{n,p}^{i,j,k}), f(m_{n,p}^{i,j,k} + 1)) \cap \dot{Y}] \geq 2,$$

which leads to a contradiction since $m_{n,p}^{i,j,k}$ has been chosen to be above i . This contradiction completes our proof. \square

Every filter \mathcal{F} gives rise to a natural forcing notion $\mathcal{M}_{\mathcal{F}}$ introducing a generic subset $X \in [\omega]^\omega$ such that $X \subset^* F$ for all $F \in \mathcal{F}$ as follows: $\mathcal{M}_{\mathcal{F}}$ consists of pairs $\langle s, F \rangle$ such that $s \in [\omega]^{<\omega}$, $F \in \mathcal{F}$, and $\max s < \min F$. A condition $\langle s, F \rangle$ is stronger than $\langle t, G \rangle$ if $F \subset G$, s is an end-extension of t , and $s \setminus t \subset G$. $\mathbb{M}_{\mathcal{F}}$ is usually called *Mathias forcing associated with \mathcal{F}* . In the proof of the next lemma we shall work with clopen subsets of $\mathcal{P}(\omega)$ of the form $\uparrow s = \{X \subset \omega : s \subset X\}$, where $s \in [\omega]^{<\omega}$.

Lemma 2.4. *Suppose that \mathcal{F} is a Hurewicz filter. Then $\mathcal{M}_{\mathcal{F}}$ is well-splitting.*

Proof. Suppose that f well-splits $M \prec H(\theta)$, and $\mathcal{F} \in M$. We shall prove that any $\langle s_0, F_0 \rangle \in \mathcal{M}_{\mathcal{F}} \cap M$ forces that f well-splits $M[\Gamma]$. This suffices because all conditions in $\mathcal{M}_{\mathcal{F}}$ are $(M, \mathcal{M}_{\mathcal{F}})$ -generic. Suppose, contrary to our claim, that there exists $\langle s_1, F_1 \rangle \leq \langle s_0, F_0 \rangle$ such that

$$\langle s_1, F_1 \rangle \Vdash \exists \sigma \exists j \exists n_0 (\sigma \in M \cap [\omega]^\omega \wedge j \in 2 \wedge n_0 \in \omega \wedge \bigwedge n \in E_j \setminus n_0 (|[f(n), f(n+1)] \cap \sigma| \leq 1)).$$

Replacing $\langle s_1, F_1 \rangle$ with a stronger condition, if necessary, we may fix $j \in 2$, $n_0 \in \omega$, and a $\mathcal{M}_{\mathcal{F}}$ -name $\dot{Y} \in M$ for an infinite subsets of ω such that

$$\langle s_1, F_1 \rangle \Vdash \forall n \in E_j \setminus n_0 (|[f(n), f(n+1)] \cap \dot{Y}| \leq 1).$$

Let $\dot{g} \in M$ be a name for a function such that $\dot{g}(n)$ is forced to be the n th element of \dot{Y} . For every $m \in \omega$ let \mathcal{S}_m be the set of those $s \in [F_0 \setminus (\max s_1 + 1)]^{<\omega}$ such that there exist $F_s \in \mathcal{F}$ such that $\langle s_1 \cup s, F_s \rangle$ forces $\dot{g}(m+1)$ to be equal to some $l_{s,m} \in \omega$. It is clear that for every $F \in \mathcal{F}$ there exists $s \in \mathcal{S}_m$ such that $s \subset F$. In other words, $\mathcal{U}_m := \{\uparrow s : s \in \mathcal{S}_m\}$ is an open cover of \mathcal{F} . Since \mathcal{F} is Hurewicz, there exists for every m a finite $\mathcal{V}_m \subset \mathcal{U}_m$ such that $\{\bigcup \mathcal{V}_m : m \in \omega\}$ is a γ -cover of \mathcal{F} . Let $\mathcal{T}_m \in [\mathcal{S}_m]^{<\omega}$ be such that $\mathcal{V}_m = \{\uparrow s : s \in \mathcal{T}_m\}$ and $h(m) = \max\{l_{s,m} : s \in \mathcal{T}_m\} + 1$. By elementarity, we can in addition assume that $\langle \mathcal{U}_m, \mathcal{V}_m, \mathcal{S}_m, \mathcal{T}_m : m \in \omega \rangle \in M$ as well as $h \in M$.

Set $h'(0) = h(0)$ and $h'(m+1) = h(h'(m))$ for all $m \in \omega$. Let m_0 be such that for every $m \geq m_0$ there exists $s \in \mathcal{S}_m \cap \mathcal{P}(F_1)$. Set $n_1 = \max\{n_0, h'(m_0)\}$. Since f well-splits M , the set $I_j := \{n \in E_j : |[f(n), f(n+1)] \cap \text{range}(h')| \geq 2\}$ is infinite, in particular it contains some $n_2 > n_1$. Thus there exists $m \in \omega$ such that

$$f(n_2) \leq h'(m) < h'(m+1) = h(h'(m)) < f(n_2).$$

f is strictly increasing, hence by the definition of n_1 we have that $m > m_0$, and therefore there exist $s \in \mathcal{S}_{h'(m)} \cap \mathcal{P}(F_1)$. Thus there exists $F_s \in \mathcal{F}$ such that

$$\langle s_1 \cup s, F_s \rangle \Vdash \dot{g}(h'(m) + 1) = l_{s, h'(m)} < h(h'(m)) < f(n_2).$$

Also, $\langle s_1 \cup s, F_s \rangle \Vdash \dot{g}(h'(m)) \geq h'(m) \geq f(n_2)$. It follows from the above that $\langle s_1 \cup s, F_s \rangle$ forces that $[f(n_2), f(n_2) + 1)$ contains at least two elements of \dot{Y} , namely the $h'(m)$ -th and $h'(m) + 1$ -st. On the other hand, $\langle s_1 \cup s, F_s \rangle$ is compatible with $\langle s_1, F_1 \rangle$ because $s \subset F_1$ and $\max s_1 < \min s$, $n_2 > n_0$, $n_2 \in E_j$, and $\langle s_1, F_1 \rangle$ forces $[f(n), f(n+1)] \cap \dot{Y} \leq 1$ for all $n \in E_j \setminus n_0$. In this way two compatible conditions $\langle s_1, F_1 \rangle$ and $\langle s_1 \cup s, F_s \rangle$ force contradictory facts, which is impossible. This completes our proof. \square

Let us mention that there is another property of posets $\mathcal{M}_{\mathcal{F}}$ for Hurewicz filters \mathcal{F} which is preserved by *finite* support iterations and which guarantees that the ground model reals remain splitting and unbounded, see [5, Prop. 84].

Corollary 2.5. *The Cohen forcing is well-splitting.*

Proof. The Cohen forcing is isomorphic to any countable atomless poset, in particular to $\mathcal{M}_{\mathfrak{F}r}$, where $\mathfrak{F}r$ is the Fréchet filter consisting of all cofinite subsets of ω . It remains to note that $\mathfrak{F}r$ is Hurewicz. \square

Recall that a poset \mathbb{P} is ω^ω -bounding if $\omega^\omega \cap V$ is dominating in $V^\mathbb{P}$.

Lemma 2.6. *Every proper ω^ω -bounding poset \mathbb{P} is well-splitting.*

Proof. Let us fix a \mathbb{P} -name \dot{Y} for an element of $[\omega]^\omega$, a countable elementary submodel M of $H(\theta)$ such that $\mathbb{P}, \dot{Y} \in M$, and $p \in \mathbb{P} \cap M$. Suppose that f well-splits M and $q \leq p$ is any (M, \mathbb{P}) -generic condition. Let $\dot{g} \in M$ be a name for the function in $\omega^{\uparrow\omega}$ which is the increasing enumeration of \dot{Y} . Since \mathbb{P} is ω^ω -bounding and q is (M, \mathbb{P}) -generic, there exist $k_0 \in \omega$ and $h \in M \cap \omega^{\uparrow\omega}$ such that $q \Vdash \dot{g}(k) < h(k)$ for all $k \geq k_0$. Let $h_1 \in M$ be the following function: $h_1(0) = 0$, $h_1(n+1) = h(h_1(n)) + 1$ for all $n \in \omega$. Let G be \mathbb{P} -generic containing q and Y, g be the evaluations of \dot{Y}, \dot{g} with respect to G , respectively. It follows from the above that the set

$$I := \{i \in E_0 : |[f(i), f(i+1)) \cap \text{range}(h_1)| \geq 2\}$$

is infinite. For every $i \in I$ we can find $n_i \in \omega$ such that $h_1(n_i), h_1(n_i + 1) \in [f(i), f(i+1))$. Thus if $i \geq k_0$ then we have

$$\begin{aligned} f(i) \leq h_1(n_i) \leq g(h_1(n_i)) &< h(h_1(n_i)) \leq g(h(h_1(n_i))) < \\ &< h(h(h_1(n_i))) = h_1(n_i + 1) < f(i), \end{aligned}$$

and hence $|[f(i), f(i+1)) \cap Y| \geq 2$ because $g(h_1(n_i)), g(h(h_1(n_i)))$ belong to the latter intersection. Therefore in $V[G]$ we have $I \subset \{i \in E_0 : |[f(i), f(i+1)) \cap Y| \geq 2\}$. Since $G \ni q$ was chosen arbitrarily, we can conclude that q forces the set

$$\{n \in E_0 : |[f(n), f(n+1)) \cap \dot{Y}| \geq 2\}$$

to be infinite, which completes our proof. \square

Summarizing the results proved in this section we get the following

Theorem 2.7. *The class of all well-splitting posets preserves ground model reals splitting and unbounded, is closed under countable support iterations, and includes ω^ω -bounding, Cohen, Miller, and Mathias forcing associated to filters with the Hurewicz covering properties.*

REFERENCES

- [1] Abraham, U., *Proper Forcing*, in: *Handbook of Set Theory* (M. Foreman, A. Kanamori, and M. Magidor, Eds.), Springer, Dordrecht 2010, pp. 333–394.
- [2] Blass, A., *Combinatorial cardinal characteristics of the continuum*, in: *Handbook of Set Theory* (M. Foreman, A. Kanamori, and M. Magidor, Eds.), Springer, Dordrecht, 2010, pp. 395–491.
- [3] Brendle, J., *Mob families and mad families*, Arch. Math. Logic **37** (1998), 183–197.
- [4] Brendle, J., *Some problems in forcing theory: large continuum and generalized cardinal invariants*, RIMS Kokyuroku 2042 (Infinite Combinatorics and Forcing Theory, T. Yorioka, ed.), 2017, 1–16.
- [5] Brendle, J.; Guzmán, O.; Raghavan, D.; Hrušák, M., *Combinatorial properties of MAD families*, preprint.

- [6] Brendle, J.; Raghavan, D., *Bounding, splitting, and almost disjointness*, Ann. Pure Appl. Logic **165** (2014), 631–651.
- [7] Chodounský, D.; Repovš, D.; Zdomsky, L., *Mathias forcing and combinatorial covering properties of filters*, J. Symb. Log. **80** (2015), 1398–1410.
- [8] Hurewicz, W., *Über Folgen stetiger Funktionen*, Fund. Math. **9** (1927), 193–204.
- [9] Just, W.; Miller, A.W.; Scheepers, M.; Szeptycki, P.J., *The combinatorics of open covers. II*, Topology Appl. **73** (1996), 241–266.
- [10] Miller, A., *Rational perfect set forcing*, in: *Axiomatic Set Theory* (J. Baumgartner, D. A. Martin, S. Shelah, Eds.), Contemporary Mathematics 31, American Mathematical Society, Providence, Rhode Island, 1984, pp. 143–159.
- [11] Shelah, S., *On cardinal invariants of the continuum*, in: *Axiomatic Set Theory* (J. Baumgartner, D. A. Martin, S. Shelah, Eds.), Contemporary Mathematics 31, American Mathematical Society, Providence, Rhode Island, 1984, pp. 183–207.
- [12] Shelah, S., *Two cardinal invariants of the continuum ($\mathfrak{d} < \mathfrak{a}$) and FS linearly ordered iterated forcing*, Acta Math. **192** (2004), 187–223.
- [13] Vaughan, J., *Small uncountable cardinals and topology*, in: *Open Problems in Topology* (J. van Mill, G.M. Reed, Eds.), Elsevier Sci. Publ., Amsterdam 1990, pp. 195–218.
- [14] Zdomsky, L., *Brendle’s proof of the consistency of $\mathfrak{b} < \mathfrak{a}$, without ranks, games, and Cohen reals*, preprint.

FACULTY OF EDUCATION, AND FACULTY OF MATHEMATICS AND PHYSICS, UNIVERSITY OF LJUBLJANA, AND INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, LJUBLJANA, SLOVENIA 1000.

E-mail address: dusan.repovs@guest.arnes.si

URL: <http://www.fmf.uni-lj.si/~repovs/index.htm>

UNIVERSITÄT WIEN, INSTITUT FÜR MATHEMATIK, KURT GÖDEL RESEARCH CENTER, AUGASSE 2–6, UZA 1 - BUILDING 2, 1090 WIEN, AUSTRIA.

E-mail address: lzdomsky@gmail.com

URL: <http://www.logic.univie.ac.at/~lzdomsky/>